

MORINGA OLEIFERA SEED DERIVATIVES
AS NATURAL BIO-COAGULANTS IN WATER
TREATMENT PROCESS

AZIZ QANNAF AZIZ ZAID

MASTER OF SCIENCE

UNIVERSITI MALAYSIA PAHANG



SUPERVISOR'S DECLARATION

We hereby declare that we have checked this thesis and in my opinion, this thesis is adequate in terms of scope and quality for the award of the degree of Master of Science.

(Supervisor's Signature)

Full Name : DR. SURIATI BINTI GHAZALI

Position :

Date :

(Co-supervisor's Signature)

Full Name : DR. NOOR SABRINA BINTI AHMAD MUTAMIM

Position :



STUDENT'S DECLARATION

I hereby declare that the work in this thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Malaysia Pahang or any other institutions.

(Student's Signature)

Full Name : AZIZ QANNAF AZIZ ZAID

ID Number : MKC17026

Date :

MORINGA OLEIFERA SEED DERIVATIVES AS NATURAL
BIO-COAGULANTS IN WATER TREATMENT PROCESS

AZIZ QANNAF AZIZ ZAID

Thesis submitted in fulfillment of the requirements
for the award of the degree of
Master of Science

Faculty of Chemical and Process Engineering Technology
UNIVERSITI MALAYSIA PAHANG

FEBRUARY 2020

ACKNOWLEDGEMENTS

All my thanks and gratitude goes to God for His protection and guidance in the course of this research work. I also acknowledge tremendous support is received from my able supervisor Dr. Suriati Binti Ghazali. I equally appreciate all the technical staffs at the Faculty of Chemical and Natural Resources Engineering for their incredible contributions during the period of my research. My thanks go to my parents and siblings for their financial support and prayers. I am much indebted to you for all your supports; May Allah grants you long life and blessings. Lastly, I appreciated my entire lab mate and all my colleagues who stood by me and assisted me, you will never lose your reward. I thank them all.

ABSTRAK

Air sungai di banyak tempat dalam dunia ini digunakan untuk tujuan minuman selepas menjalani beberapa rawatan. Akibatnya, kekeruhan dalam air yang tidak dirawat secara konvensional dikeluarkan dengan merawat air melalui bahan kimia mahal, yang kebanyakannya tidak mesra alam. Malah, dengan menggunakan bahan-bahan kimia dalam rawatan air mempunyai kesan negatif kepada kesihatan manusia. Oleh itu, terdapat keperluan untuk menggunakan bahan semula jadi sebagai alternatif untuk menggantikan bahan kimia. *Moringa oleifera* adalah tumbuhan komoditi penting yang telah digunakan untuk rawatan air di bahagian tropika di dunia. Dalam kajian ini, potensi benih oleifera *Moringa* sebagai bio-koagulan untuk rawatan air telah disiasat pada keadaan optimum. Eksperimen ini dijalankan dalam dua peringkat, kaedah yang pertama telah digunakan untuk menentukan saiz zarah terbaik dan jumlah *Moringa oleifera* benih serbuk (*MOSP*) untuk rawatan air. Ia adalah diperhatikan bahawa saiz zarah 2 mm prestasi yang lebih baik dari segi kecekapan kekeruhan penyingkiran 4.98 NTU, dengan 0.1 g. Tambahan pula, kaedah *MOSP* kedua diletakkan dalam pengekstrakan bidal untuk mengeluarkan minyak dari benih dan dihasilkan *MOCR* digunakan sebagai koagulan biologi dalam rawatan air. Kaedah gerak balas permukaan telah digunakan untuk menentukan kesan parametrik mencampurkan kelajuan, masa dan dos pada kekeruhan sisa. Ketiga-tiga faktor, seperti kelajuan yang rendah, masa kelajuan rendah dan dos disiasat menggunakan Metodologi Response Surface (RSM) untuk optimum penyingkiran kekeruhan air. Keputusan yang diperolehi daripada reka bentuk eksperimen menunjukkan keadaan optimum pada kelajuan yang rendah, masa kelajuan rendah dan dos 40 rpm, 60 min, dan 0.75 mg / L, masing-masing. Dalam siasatan ini meramalkan (teori) sisa kekeruhan adalah 4.73 NTU. Hasil daripada ANOVA untuk pengoptimuman kekeruhan sisa menunjukkan bahawa model kuadratik adalah signifikan pada tahap keyakinan 95% ($p < 0.05$). Perbezaan antara kekeruhan teori dan eksperimen menunjukkan bahawa tiada perbezaan signifikan dengan ralat peratusan 1.6913%. Kualiti air yang dirawat telah ditentukan pada keadaan optimum menggunakan ujian kualitatif. Hasil penentuan kualiti air menunjukkan bahawa kekeruhan, kekonduksian, TDS, COD dan BOD dengan masing-masing menunjukkan peratusan sebanyak 99%, 73%, 50%, 75% dan 28%. Akhirnya, keputusan yang diperolehi menunjukkan potensi *Moringa oleifera* sebagai koagulan semula jadi dalam rawatan berkesan air untuk tujuan minum. Kekeruhan yang lebih rendah (< 5 NTU) yang dicapai daripada kajian ini mengesahkan penggunaan produk semulajadi yang mesra alam yang penting ini untuk rawatan air.

ABSTRACT

River water in many parts of the world used for drinking purpose after some treatments. Consequently, the turbidity in untreated water is often conventionally removed by treating the water with expensive chemicals, many of which are not environmentally friendly. In fact, using chemicals materials in water treatments have some negative effects to human health. Therefore, there is a need to use some natural materials as an alternative to replace chemicals. *Moringa oleifera* is an important commodity plants which has been used for the treatment of water in tropical part of the world. In this study, the potential of *Moringa oleifera* seeds as bio-coagulant for water treatment were investigated at optimum condition. The experiment was conducted in two stages, the first method of was employed to determine which of the best particle sizes and amount of *Moringa oleifera* seeds powder (*MOSP*) for water treatment. It was observed that the 2 mm particle size performed better in terms of turbidity removal efficiency 4.98NTU, with 0.1 g. Furthermore, the second method *MOSP* was placed in extraction thimble to remove the oil from the seeds and produced *MOCR* used as bio- coagulant in water treatment. Response surface methodology was applied to determine the parametric effects of mixing speed, time and dosage on the residual turbidity. The three independent factors, such as Low speed, Low speed time and dosage were optimized using the Response Surface Methodology (RSM) for optimal water turbidity removal. The results obtained from experimental design demonstrated the optimum condition at low speed, low speed time and dosage as 40 rpm, 60 min, and 0.75 mg/L, respectively. Under this condition the predicted (theoretical) residual turbidity was 4.73 NTU. The result of the ANOVA for the optimization of the residual turbidity showed that the quadratic model was significant at 95% confidence level ($p < 0.05$). The difference between the theoretical and experimental turbidity showed that no significant variation with a percentage error value of 1.6913 %. The quality of the treated water was determined at optimum condition using qualitative tests. The result of the water quality determination revealed that turbidity, conductivity, TDS, COD and BOD with corresponding percentage changes of 99 %, 73 %, 50 %, 75 % and 28 %, respectively. Finally, the result obtained therefore showed the potential of *Moringa oleifera* as natural coagulants in the effective treatment of water for drinking purpose. The lower turbidity (<5 NTU) achieved from this study confirmed the potential of this important eco-friendly natural product for the treatment of water.

TABLE OF CONTENT

DECLARATION	
TITLE PAGE	
ACKNOWLEDGEMENTS	ii
ABSTRAK	iii
ABSTRACT	iv
TABLE OF CONTENT	v
LIST OF TABLES	ix
LIST OF FIGURES	x
CHAPTER 1 INTRODUCTION	1
1.1 Introduction	1
1.2 Problem Statement	2
1.3 Research Objectives	3
1.4 Research Scopes	4
1.5 Significance of Research	4
CHAPTER 2 LITERATURE REVIEW	6
2.1 Introduction	6
2.2 Water Treatment	7
2.3 Coagulation – flocculation in water and wastewater treatment	9
2.3.1 Mechanism of Coagulation Process	9
2.3.2 Types of Coagulants	10
2.4 <i>Moringa oleifera</i>	16
2.4.1 Application of <i>Moringa Oleifera</i> Seeds in Water Treatment	18
2.4.2 <i>Moringa Oleifera</i> Seeds Oil Extraction	22

2.4.3	Active Component Extraction from (<i>MOCR</i>).	23
2.5	Optimization of coagulation using RSM	23
2.6	Water Quality	24
2.6.1	Turbidity	24
2.6.2	Biological Oxygen Demand (BOD)	25
2.6.3	Chemical Oxygen Demand (COD)	26
2.6.4	pH	26
2.6.5	Conductivity and Total Dissolved Solids (TDS)	27
2.6.6	Overall Water Quality Performance	28
CHAPTER 3 MATERIALS AND METHODOLOGY		30
3.1	Introduction	30
3.2	Sample Preparation	32
3.2.1	<i>Moringa oleifera</i> Seeds Powder (MOSP)	32
3.2.2	Preparation of <i>Moringa oleifera</i> Cake Residue (MOCR) through extraction process	34
3.2.3	Salt extraction	35
3.3	Water sample Collection	36
3.4	Design of Experiment to optimize the study	36
3.5	Jar Test	37
3.6	Water Quality Analysis	37
3.6.1	Turbidity Measurement	37
3.6.2	Chemical Oxygen Demand (COD) Measurement	38
3.6.3	Biochemical Oxygen Demand (BOD) Measurement	40
3.6.4	pH, Conductivity and Total Dissolved Solid Measurement	40

CHAPTER 4 RESULTS AND DISCUSSION	42
4.1 Introduction	42
4.2 Characterizations of Belat River water	42
4.3 Determination <i>MOSP</i> for Water Treatment	43
4.3.1 Turbidity	43
4.3.2 pH	44
4.3.3 Conductivity	45
4.3.4 TDS Measurement	46
4.3.5 COD Measurement	47
4.3.6 BOD Measurement	48
4.4 <i>Moringa oleifera</i> Cake Recovery (MOCR)	49
4.5 Optimization Study of Operating Parameters in Water Treatment Using (RSM).	50
4.6 Model Fitting and Analysis of Variance	51
4.6.1 Validation of Predictive Model	53
4.6.2 The Effects of Variables on the Optimal Settings	55
4.7 Quality Analysis of Treated Water	56
4.7.1 pH Measurement	56
4.7.2 Total Dissolved Solid (TDS)	57
4.7.3 Conductivity	58
4.7.4 COD Measurement	59
4.7.5 BOD Measurement	60
4.7.6 Summary of Water Quality Analysis	62
CHAPTER 5 CONCLUSION AND RECOMMENDATION	63
5.1 Overall Conclusions	63
5.2 Recommendation	64

REFERENCES	65
APPENDICES	75
APPENDIX A RESULTS OF QUALITY CHARACTERISTICS MEASUREMENT (MOSP)	75
APPENDIX B RESULTS OF QUALITY CHARACTERISTICS MEASUREMENT (MO BIO-COAGULANT)	78

LIST OF TABLES

Table 2.1	Advantages and disadvantages of some methods for water treatment	8
Table 2.2	Uses of various part from <i>Moringa oleifera</i>	17
Table 2.3	Summary of previous research on <i>Moringa oleifera</i> and their significant information and results.	19
Table 2.4	Percentages of Oil Extracted with Three Different Solvents	22
Table 2.5	Ministry of Health Malaysia (2014)	28
Table 2.6	National Water Quality Standards for Malaysia (WEPA, 2017)	28
Table 2.7	National Water Quality Classes for Malaysia (WEPA, 2017)	29
Table 3.1	Coded and actual design range	37
Table 4.1	Summary of result for water quality characteristics	42
Table 4.2	Design range, experimental design and results obtained from optimization	51
Table 4.3	Analysis of variance for response surface quadratic model	52
Table 4.4	Theoretical optimal conditions	54
Table 4.5	Validation of experimental best condition	54
Table 4.6	Summary of result for water quality characteristics Using <i>Moringa oleifera</i> bio-coagulant	62

LIST OF FIGURES

Figure 2-1	Drinking water treatment plant	7
Figure 2-2	<i>Parkinsonia aculeate</i>	12
Figure 2-3	<i>Cicer arietinum</i> and <i>Dolichos lablab</i>	13
Figure 2-4	<i>Cassia alata</i> Leaves	14
Figure 2-5	Peanut seeds	15
Figure 2-6	<i>Jatropha curcas</i> seeds	16
Figure 2-7	<i>Moringa oleifera</i> Pods and Seeds	17
Figure 2-8	pH Value Range	27
Figure 3-1	Process Flow Chart of Experiment	31
Figure 3-2	Undehusk MO seeds and Dehusk MO seeds	32
Figure 3-3	Domestic grinder to grinding MO	33
Figure 3-4	Electronic sieve shaker	33
Figure 3-5	MOSP after sieved and packed into clean, polythene bags and sealed	33
Figure 3-6	Soxhlet Extractor	34
Figure 3-7	Salt extraction	35
Figure 3-8	River water sample collection	36
Figure 3-9	Paddle Jar Test Apparatus	37
Figure 3-10	Digital turbidity meter (TB-500G) and (DTC-4DG)	38
Figure 3-11	COD rack and COD reactor HACH DRB200	39
Figure 3-12	HACH spectrophotometer DR2800	39
Figure 3-13	Manual Calibration for pH before measurement	41
Figure 3-14	pH measurement for the mixture	41
Figure 4-1	Turbidity removal of water treated with <i>MOSP</i>	43
Figure 4-2	pH values for water treated with MOSP	45
Figure 4-3	Water conductivity treated with MOSP	46
Figure 4-4	TDS of water treated with MOSP	47
Figure 4-5	COD of water treated with MOSP	48
Figure 4-6	BOD of water treated with MOSP	49
Figure 4-7	The interaction plot (a) low speed and low speed mixing time (b) Dosage and low speed (c) Dosage and low speed time	53
Figure 4-8	pH Values after Treatment	57
Figure 4-9	TDS Values after Treatment	58
Figure 4-10	Conductivity Values after Treatment	59

Figure 4-11	COD Values after Treatment	60
Figure 4-12	BOD Values after Treatment	61

REFERENCES

- Abdalla, Elsadig Eshag, Yousif Ali Yousif, and Ibnaof Ali Ibnaof. "Comparative study between Aluminium sulphate and Moringa Oleifera seeds as coagulants in drinking water treatment." *University Of Khartoum Engineering Journal* 8, no. 1 (2018).
- Abiola, O. N. (2019). *Polymers for Coagulation and Flocculation in Water Treatment*. In *Polymeric Materials for Clean Water* (pp. 77-92). Springer, Cham.
- Adeniran, K. A., Akpenpuun, T. D., Akinyemi, B. A., & Wasiu, R. A. (2017). Effectiveness of Moringa oleifera seed as a coagulant in domestic wastewater treatment. *African Journal of Science, Technology, Innovation and Development*, 9(3), 323-328
- Ahmad, T., Ahmad, K., & Alam, M. (2016). Sustainable management of water treatment sludge through 3 'R' concept. *Journal of Cleaner Production*, 124, 1-13.
- Alfarra, R.S., Ali, E.N. and Yusoff, M.M. (2014). Removal of Heavy Metals by Natural Adsorbent. *International Journal of Biosciences*, 4(7), 130-139.
- Ali, Eman N., Sabreen R. Alfarra, Mashita Mohd Yusoff, and Md Lutfor Rahman. "Environmentally friendly biosorbent from Moringa oleifera leaves for water treatment." *International Journal of Environmental Science and Development* 6, no. 3 (2015): 165.
- Ali, E. N., Muyibi, S. A., Alam, M. Z., & Salleh, H. M. (2012, September). Optimization of water treatment parameters using processed Moringa oleifera as a natural coagulant for low turbidity water. In *Statistics in Science, Business, and Engineering (ICSSBE), 2012 International Conference on* (pp. 1-6). IEEE.
- Ali, E. N., Muyibi, S. A., Salleh, H. M., Alam, M. Z., & Salleh, M. R. M. (2010). Production of natural coagulant from Moringa oleifera seed for application in treatment of low turbidity water. *Journal of Water Resource and Protection*, 2(03), 259.
- Ali, E. N., Muyibi, S. A., Salleh, H. M., Salleh, M. R. M., & Alam, M. Z. (2009). Moringa oleifera seeds as natural coagulant for water treatment. In *Thirteenth International Water Technology Conference, IWTC* (Vol. 13, No. 2009, pp. 163-168).
- Al-Kalifawi, E. J. (2016). Purification of the Tigris River water using seeds extract of Al-Ruwag tree (*Moringa oleifera* Lamarck) cultivated in Iraq. *Magazine of Al-Kufa University for Biology, (Special Second International Scientific Conference for the Life Sciences)*, 14-22.
- Al-Mamun, A., & Zainuddin, Z. (2013). Sustainable river water quality management in Malaysia. *IIUM Engineering Journal*, 14(1).

- Anjorin, T. S., Ikokoh, P., & Okolo, S. (2010). Mineral composition of *Moringa oleifera* leaves, pods and seeds from two regions in Abuja, Nigeria. *International Journal of Agriculture and Biology*, 12(3), 431-434.
- Anwar, F., Latif, S., Ashraf, M. and Gilani, A.H. (2007). *Moringa oleifera*: A Food Plant with Multiple Medicinal Uses. *Phytotherapy Research*, 21, 17-25.
- Anwar, F., & Rashid, U. (2007). Physico-chemical characteristics of *Moringa oleifera* seeds and seed oil from a wild provenance of Pakistan. *Pak. J. Bot*, 39(5), 1443-1453.
- Amagloh, F. K., & Benang, A. (2009). Effectiveness of *Moringa oleifera* seed as coagulant for water purification. *African Journal of Agricultural Research*, 4(2), 119-123.
- Arnoldsson, E., Bergman, M., & Matsinhe, N. (2008). Assessment of drinking water bark extracts of *Moringa oleifera* in reducing bacterial load in water. *Int J Adv Res*, 4, 124-130.
- Baskan, M. B., & Pala, A. (2010). A statistical experiment design approach for arsenic removal by coagulation process using aluminum sulfate. *Desalination*, 254(1-3), 42-48.
- Batool, A., Ziaf, K., & Amjad, M. (2015). Effect of halo-priming on germination and vigor index of cabbage (*Brassica oleracea* var. capitata). *Journal of Environmental and Agricultural Sciences*, 2(7), 8pp.
- Bezerra, M. A., Santelli, R. E., Oliveira, E. P., Villar, L. S., & Escalera, L. A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 76(5), 965-977.
- Birima, A. H., Hammad, H. A., Desa, M. N. M., & Muda, Z. C. (2013). Extraction of natural coagulant from peanut seeds for treatment of turbid water. In *IOP Conference Series: Earth and Environmental Science* (Vol. 16, No. 1, p. 012065). IOP Publishing.
- Bodlund, I. (2013). Coagulant protein from plant materials: Potential water treatment agent (Doctoral dissertation, KTH Royal Institute of Technology).
- Bookter, A. (2009). *Water Quality Trends in the Entiat River Sub-basin: 2007-2008* (Vol. 563). DIANE Publishing.
- Bratby, J. (2016). *Coagulation and flocculation in water and wastewater treatment*. IWA publishing.
- Brunet, A., Tardin, C., Salome, L., Rousseau, P., Destainville, N., & Manghi, M. (2015). Dependence of DNA persistence length on ionic strength of solutions with monovalent and divalent salts: a joint theory–experiment study. *Macromolecules*, 48(11), 3641-3652.

- Brusseau, M. L., & Artiola, J. F. (2019). Chemical contaminants. In *Environmental and pollution science* (pp. 175-190). Academic Press.
- Byrne, M. D. (2013, July). How many times should a stochastic model be run? An approach based on confidence intervals. In *Proceedings of the 12th International conference on cognitive modeling*, Ottawa.
- Chamarro, E., Marco, A., & Esplugas, S. (2001). Use of Fenton reagent to improve organic chemical biodegradability. *Water research*, 35(4), 1047-1051.
- Chattoraj, S., Mondal, N. K., & Sen, K. (2018). Removal of carbaryl insecticide from aqueous solution using eggshell powder: a modeling study. *Applied Water Science*, 8(6), 163.
- Chen, J., Liu, S., Qi, X., Yan, S., & Guo, Q. (2018). Study and design on chemical oxygen demand measurement based on ultraviolet absorption. *Sensors and Actuators B: Chemical*, 254, 778-784.
- Chen, S., Yuan, Z., Hanigan, D., Westerhoff, P., Zhao, H., & Ni, J. (2018). Coagulation behaviors of new covalently bound hybrid coagulants (CBHyC) in surface water treatment. *Separation and Purification Technology*, 192, 322-328.
- Cheremisinoff, N. P. (2001). *Handbook of water and wastewater treatment technologies*. Butterworth-Heinemann.
- Chew, C. M., Aroua, M. K., Hussain, M. A., & Ismail, W. M. Z. W. (2016). Evaluation of ultrafiltration and conventional water treatment systems for sustainable development: an industrial scale case study. *Journal of cleaner production*, 112, 3152-3163.
- Choi, Y., Yang, T. D., Fang-Yen, C., Kang, P., Lee, K. J., Dasari, R & Choi, W. (2011). Overcoming the diffraction limit using multiple light scattering in a highly disordered medium. *Physical review letters*, 107(2), 023902.
- Choy, S. P. (2015). A review on common vegetables and legumes as promising plant-based natural coagulants in water clarification. *International Journal of Environmental Science and Technology*, 12 (1), 367–390.
- Chupin, L., Maunu, S. L., Reynaud, S., Pizzi, a, Charrier, B., & Bouhtoury, F. C.-E. (2015). Microwave assisted extraction of maritime pine (*Pinus pinaster*) bark: Impact of particle size and characterization. *Industrial Crops and Products*, 65(1), 142–149.
- Da Porto, C., Decorti, D., & Natolino, A. (2016). Microwave pretreatment of *Moringa oleifera* seed: effect on oil obtained by pilot-scale supercritical carbon dioxide extraction and Soxhlet apparatus. *The Journal of Supercritical Fluids*, 107, 38-43.
- Dehghani, M., & Alizadeh, M. H. (2016). The effects of the natural coagulant *Moringa oleifera* and alum in wastewater treatment at the Bandar Abbas Oil Refinery. *Environmental Health Engineering and Management Journal*, 3(4), 225-230.

- Delzer, G. C., & McKenzie, S. W. (2003). Five-day biochemical oxygen demand. US Geological Survey Techniques of Water-Resources Investigations, 3.
- Demirbas, E., & Kobya, M. (2017). Operating cost and treatment of metalworking fluid wastewater by chemical coagulation and electrocoagulation processes. *Process Safety and Environmental Protection*, 105, 79-90.
- Dhanorkar, S., & Kamra, A. K. (2001). Effect of coagulation on the particle charge distribution and air conductivity. *Journal of Geophysical Research: Atmospheres*, 106(D11), 12055-12065.
- Eman, N. A., Tan, C. S., & Makky, E. A. (2014). Impact of Moringa oleifera cake residue application on waste water treatment: a case study. *Journal of Water Resource and Protection*, 6(7), 677.
- Emmanuel, S. A., Zaku, S. G., Adedirin, S. O., Tafida, M., & Thomas, S. A. (2011). Moringa oleifera seed-cake, alternative biodegradable and biocompatibility organic fertilizer for modern farming. *Magnesium*, 203, 0-08.
- Farah, N., & Torell, G. (2018). Defensive Investment in Municipal Water Hardness Reduction. Available at SSRN 3279633.
- Farooq, U., Kozinski, J. A., Khan, M. A., & Athar, M. (2010). Biosorption of heavy metal ions using wheat based biosorbents—a review of the recent literature. *Bioresource technology*, 101(14), 5043-5053.
- Fletcher, S. M., Stark, D., Harkness, J., & Ellis, J. (2012). Enteric protozoa in the developed world: a public health perspective. *Clinical microbiology reviews*, 25(3), 420-449.
- Foidl, N., Makkar, H. P. S., & Becker, K. (2001). The potential of Moringa oleifera for agricultural and industrial uses. *The miracle tree: The multiple attributes of Moringa*, 45-76.
- Fu, F., & Wang, Q. (2011). Removal of heavy metal ions from wastewaters: a review. *Journal of environmental management*, 92(3), 407-418.
- Gelda, R. K., Effler, S. W., Peng, F., Owens, E. M., & Pierson, D. C. (2009). Turbidity model for ashokan reservoir, New York: case study. *Journal of Environmental Engineering*, 135(9), 885-895.
- Ghaly, A. E., Ananthashankar, R., Alhattab, M. V. V. R., & Ramakrishnan, V. V. (2014). Production, characterization and treatment of textile effluents: a critical review. *J Chem Eng Process Technol*, 5(1), 1-19.
- Goudeli, E., Eggersdorfer, M. L., & Pratsinis, S. E. (2015). Coagulation–Agglomeration of fractal-like particles: Structure and self-preserving size distribution. *Langmuir*, 31(4), 1320-1327.

- Green, D. A., Browning, G. A., & Schaeffer, D. L. (2018). U.S. Patent Application No. 15/433,652.
- Harisha, R. S., Hosamani, K. M., Keri, R. S., Nataraj, S. K., & Aminabhavi, T. M. (2010). Arsenic removal from drinking water using thin film composite nanofiltration membrane. *Desalination*, 252(1-3), 75-80
- Hendrawati, Yuliastri, I. R., Nurhasni, Rohaeti, E., Effendi, H., & Darusman, L. K. (2016). The use of *Moringa Oleifera* Seed Powder as Coagulant to Improve the Quality of Wastewater and Ground Water. *IOP Conference Series: Earth and Environmental Science*, 31, 12033.
- Hicks, B. M., South, S. C., DiRago, A. C., Iacono, W. G., & McGue, M. (2009). Environmental adversity and increasing genetic risk for externalizing disorders. *Archives of general psychiatry*, 66(6), 640-648.
- Hoa, N. T., & Hue, C. T. (2018). Enhanced water treatment by *Moringa oleifera* seeds extract as the bio-coagulant: role of the extraction method. *Journal of Water Supply: Research and Technology-Aqua*, 67(7), 634-647
- Hooda, P. S., Edwards, A. C., Anderson, H. A., & Miller, A. (2000). A review of water quality concerns in livestock farming areas. *Science of the Total Environment*, 250(1-3), 143-167.
- Idris, M. A., Jami, M. S., Hammed, A. M., & Jamal, P. (2016). *Moringa oleifera* seed extract: A review on its environmental applications. *Int. J. Appl. Environ. Sci*, 11, 1469-1486.
- Ikonomopoulou, M. P., Olszowy, H., Limpus, C., Francis, R., & Whittier, J. (2011). Trace element concentrations in nesting flatback turtles (*Natator depressus*) from Curtis Island, Queensland, Australia. *Marine Environmental Research*, 71(1), 10-16.
- Ju, J., Liu, R., He, Z., Liu, H., Zhang, X., & Qu, J. (2016). Utilization of aluminum hydroxide waste generated in fluoride adsorption and coagulation processes for adsorptive removal of cadmium ion. *Frontiers of Environmental Science & Engineering*, 10(3), 467-476.
- Kalibbala, H. M. (2007). Application of indigenous materials in drinking water treatment (Doctoral dissertation, KTH Royal Institute of Technology).
- Karri, V. R., Samiwala, M., & Kothapalli, L. (2016). Effect of powdered *Moringa oleifera* seeds on the characteristics of surface water. *Indian Journal of Applied Research*, 8.
- Kasmawati, M. L. (2007). *Methods of Analysis for Water and Wastewater*. UiTM, Shah Alam.
- Kemp, P., Sear, D., Collins, A., Naden, P., & Jones, I. (2011). The impacts of fine sediment on riverine fish. *Hydrological Processes*, 25(11), 1800-1821.

- Kim, B. H., Chang, I. S., Gil, G. C., Park, H. S., & Kim, H. J. (2003). Novel BOD (biological oxygen demand) sensor using mediator-less microbial fuel cell. *Biotechnology letters*, 25(7), 541-545.
- Lachin, K., Le Sauze, N., Raimondi, N. D. M., Aubin, J., Cabassud, M., & Gourdon, C. (2019). Estimation of characteristic coagulation time based on brownian coagulation theory and stability ratio modeling using electrokinetic measurements. *Chemical Engineering Journal*.
- Latifah, H. E. R. H. E., & Darusman, K. (2015). Characterization of Physico-Chemical Properties of Nano-Sized Moringa oleifera Seed Powder and Its Application as Natural Coagulant in Water Purification Process. 5 (21), 19-26.
- Lawrence, M. B., Joseph, J., Phondekar, K., & Moodi, K. (2019). DC conductivity behaviour of poly (vinyl alcohol)-based ferrogels: role of borax and carbonyl iron. *Polymer Bulletin*, 1-15.
- Levallois, P., & Villanueva, C. M. (2019). Drinking Water Quality and Human Health: An Editorial.
- Nasirian, M. (2007). A new water quality index for environmental contamination contributed by mineral processing: a case study of Amang (Tin Tailing) processing activity. *Journal of applied sciences*, 7(20), 2977-2987.
- Neethu, B., & Ghangrekar, M. M. (2017). Electricity generation through a photo sediment microbial fuel cell using algae at the cathode. *Water Science and Technology*, 76(12), 3269-3277.
- Nordmark, B. A., Bechtel, T. M., Riley, J. K., Velegol, D., Velegol, S. B., Przybycien, T. M., & Tilton, R. D. (2018). Moringa oleifera seed protein adsorption to silica: Effects of water hardness, fractionation, and fatty acid extraction. *Langmuir*, 34(16), 4852-4860.
- Nouhi, S., Kwaambwa, H. M., Gutfreund, P., & Rennie, A. R. (2018). Comparative study of flocculation and adsorption behavior of Moringa peregrina and Moringa oleifera seed proteins for water treatment.
- Nouman, Wasif, Shahzad Maqsood Ahmed Basra, Muhammad Tahir Siddiqui, Azra Yasmeen, Tehseen Gull, and Maria Angelica Cervantes Alcayde. "Potential of Moringa oleifera L. as livestock fodder crop: a review." *Turkish Journal of Agriculture and Forestry* 38, no. 1 (2014): 1-14.
- Madzvamuse, A. (2015). A Study of The Removal of Pb, Cd, and Cr Heavy Metals from Aqueous Solutions by Moringa oleifera Seeds in Comparison to Carbonized Moringa oleifera Seeds. *University of Zimbabwe*, 1-59.
- Maizuwo, A. I., Hassan, A. S., Momoh, H., & Muhammad, J. A. (2017). Phytochemical constituents, biological activities, therapeutic potentials and nutritional values of Moringa oleifera (Zogale): a review. *Journal of drug design and medicinal chemistry*, 3(4), 60.

- McConnachie, G. L., Folkard, G. K., Mtawali, M. A., & Sutherland, J. P. (1999). Field trials of appropriate hydraulic flocculation processes1. *Water Research*, 33(6), 1425-1434.
- Menezes, J. C. C. C., Colling, A. V., Silva, R. A. S., Dos Santos, R. H., & Scheneider, I. A. H. (2017). Ferric Sulphate Coagulant Obtained by Leaching from Coal Tailings. *Mine Water and the Environment*, 36(3), 457-460.
- Miraji. H. (2014). Performance Comparison between Moringa Oleifera and Alum in Water and Wastewater Treatment: A Non-synergic approach. *International Journal of Current Research*, 9198-9200.
- Monmaturapoj, N. (2017). Nano-size hydroxyapatite powders preparation by wet-chemical precipitation route. *Journal of Metals, Materials and Minerals*, 18(1).
- Ogunsina, B. S., Indira, T. N., Bhatnagar, A. S., Radha, C., Debnath, S., & Krishna, A. G. (2014). Quality characteristics and stability of Moringa oleifera seed oil of Indian origin. *Journal of food science and technology*, 51(3), 503-510.
- Okuda, T., Baes, A. U., Nishijima, W., & Okada, M. (1999). Improvement of extraction method of coagulation active components from Moringa oleifera seed. *Water research*, 33(15), 3373-3378.
- Onoji, S. E., Iyuke, S. E., Igbafe, A. I., & Nkazi, D. B. (2016). Rubber seed oil: A potential renewable source of biodiesel for sustainable development in sub-Saharan Africa. *Energy conversion and management*, 110, 125-134.
- Oram B. (2014). The pH of Water. Retrieved from Water Research Centre: <http://www.water-research.net/index.php/ph>.
- Patil, A., DasGupta, A., & Eriksson, A. (2015). Contact mechanics of a circular membrane inflated against a deformable substrate. *International Journal of Solids and Structures*, 67, 250-262.
- Paula, H.M., Oliveira Ilha, M.S. and Andrade, L.S. (2014). Concrete Plant Wastewater Treatment Process by Coagulation Combining Aluminum Sulfate and Moringa oleifera Powder. *Journal of Cleaner Production*, 76, 125-130.
- Peng, Tey Nai, and Lai Siow Li. "POPULATION GROWTH AND DISTRIBUTION IN MALAYSIA." Kuala Lumpur, Malaysia 2019 (2019): 32.
- Pontual, E. V., de Lima Santos, N. D., de Moura, M. C., Coelho, L. C. B. B., Navarro, D. M. D. A. F., Napoleão, T. H., & Paiva, P. M. G. (2014). Trypsin inhibitor from Moringa oleifera flowers interferes with survival and development of Aedes aegypti larvae and kills bacteria inhabitant of larvae midgut. *Parasitology research*, 113(2), 727-733.
- Prasad, S.V.M. and Rao, B.S. (2016). Influence of Plant-Based Coagulants in Waste Water Treatment. *International Journal of Latest Technology in Engineering, Management & Applied Science*, 5(3), 45-48.

- Rak, A. E., & Ismail, A. A. M. (2012). Cassia alata as a potential coagulant in water treatment. Research journal of recent sciences, 1(2), 28-33.
- Rafferty, K. (1999). Scaling in geothermal heat pump systems. Klamath Falls, OR: Geo-Heat Center.
- Ravikumar, K. and Sheeja, A.K. (2013). Heavy Metal Removal from Water using Moringa oleifera Seed Coagulant and Double Filtration. International Journal of Scientific & Engineering Research, 4(5), 10-13.
- Robson, C., & McCartan, K. (2016). Real world research. John Wiley & Sons.
- Rubio, J., Souza, M. L., & Smith, R. W. (2002). Overview of flotation as a wastewater treatment technique. Minerals Engineering, 15(3), 139-155.
- Sancho et al. (2016). Moringa oleifera for drinking water treatment: influence of the solvent and method used in oil-extraction on the coagulant efficiency of the seed extract. Desalination and Water Treatment, 48 - 49.
- Saritha, e. a. (2017). A Review on Native Plant Based Coagulants for Water Purification. International Journal of Applied Environmental Sciences, 18. Retrieved 3 October, 2017, from <http://www.ripublication.com>
- Sasikala S, Muthuraman G. (2016) A Laboratory Study for the Treatment of Turbidity and Total Hardness Bearing Synthetic Wastewater/Ground Water Using Moringa oleifera. Ind Chem 2:112.
- Seidi, S., Yamini, Y., Baheri, T., & Feizbakhsh, R. (2011). Electrokinetic extraction on artificial liquid membranes of amphetamine-type stimulants from urine samples followed by high performance liquid chromatography analysis. Journal of Chromatography A, 1218(26), 3958-3965.
- Shan, T. C., Al Matar, M., Makky, E. A., & Ali, E. N. (2017). The use of Moringa oleifera seed as a natural coagulant for wastewater treatment and heavy metals removal. Applied Water Science, 7(3), 1369-1376.
- Sharma, P., Kumari, P., Srivastava, M.M. & Srivastava, S. (2006). Removal of cadmium from aqueous system by shelled Moringa oleifera Lam. seed powder, 97, 299–305.
- Siddhuraju, P., & Becker, K. (2003). Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (Moringa oleifera Lam.) leaves. Journal of agricultural and food chemistry, 51(8), 2144-2155.
- Singh, R. (2014). Membrane technology and engineering for water purification: application, systems design and operation. Butterworth-Heinemann.
- Sohrabi, Y., Rahimi, S., Nafez, A. H., Mirzaei, N., Bagheri, A., Kamal Ghadiri, S & Charganeh, S. S. (2018). Chemical coagulation efficiency in removal of water turbidity. International Journal of Pharmaceutical Research, 10(3).

- Srikanth, M. V., Sunil, S. A., Rao, N. S., Uhumwangho, M. U., & Murthy, K. R. (2010). Ion-exchange resins as controlled drug delivery carriers. *Journal of Scientific Research*, 2(3), 597.
- Suhartini, S. H. (2013). Influence of Powdered Moringa oleifera Seeds and Natural Filter Media on the Characteristics of Tapioca Starch Wastewater. *International Journal of Recycling of Organic Waste in Agriculture*, 2, 1-11.
- Swistock, B. R., Clemens, S., Sharpe, W. E., & Rummel, S. (2013). Water quality and management of private drinking water wells in Pennsylvania. *Journal of environmental health*, 75(6), 60-67.
- Tariq, M., Ahmad, M., Siddique, S., Waheed, A., Shafiq, T., & Khan, M. H. (2012). Optimization of coagulation process for the treatment of the characterized slaughterhouse wastewater. *Pakistan Journal of Scientific & Industrial Research Series A: Physical Sciences*, 55(1), 43-48.
- Thatai, S., Verma, R., Khurana, P., Goel, P., & Kumar, D. (2019). Water quality standards, its pollution and treatment methods. In *A New Generation Material Graphene: Applications in Water Technology* (pp. 21-42). Springer, Cham.
- Thomas, P. A., Kern, D. E., Hughes, M. T., & Chen, B. Y. (Eds.). (2016). *Curriculum development for medical education: a six-step approach*. JHU Press.
- Tunggolou, J., & Payus, C. (2017). Moringa oleifera as coagulant used in water purification process for consumption. *Earth Sci. Pak*, 1, 1-3.
- Tzoupanos, N. D., & Zouboulis, A. I. (2008, August). Coagulation-flocculation processes in water/wastewater treatment: the application of new generation of chemical reagents. In *6th IASME/WSEAS International Conference Greece*.
- Verma, D., Fortunati, E., Jain, S., & Zhang, X. (2019). *Biomass, Biopolymer-Based Materials, and Bioenergy: Construction, Biomedical, and other Industrial Applications*.
- Wei, Z., Hou, J., & Zhu, Z. (2016). High-aluminum fly ash recycling for fabrication of cost-effective ceramic membrane supports. *Journal of Alloys and Compounds*, 683, 474-480.
- Yongabi, K. A. (2012). *A Sustainable Low-Cost Phytodisinfectant- Sand Filter Alternative for Water Purification*. The University of Adelaide Australia, 1-207.
- Yusuf, J., Yuakubu, M. B., & Balarabe, A. M. (2015). The Use of Moringa Oleifera Seed As A Coagulant For Domestic Water Purification. *IOSR Journal of Pharmacy and Biological Sciences Ver. III*, 10(1), 2319–7676.
- Yu, W., Gregory, J., & Campos, L. (2010). The effect of additional coagulant on the re-growth of alum–kaolin flocs. *Separation and Purification Technology*, 74(3), 305-309.

Zhu, G., Zheng, H., Chen, W., Fan, W., Zhang, P., & Tshukudu, T. (2012). Preparation of a composite coagulant: Polymeric aluminum ferric sulfate (PAFS) for wastewater treatment. *Desalination*, 285, 315-323.